

## **Cheap DECAF: Density Estimation for Cetaceans from Acoustic Fixed Sensors Using Separate, Non-Linked Devices**

Len Thomas  
Centre for Research into Ecological and Environmental Modelling (CREEM)  
University of St Andrews  
St Andrews, UK.  
phone: UK+1334 461801 fax: UK+1334 461800 email: [len@mcs.st-and.ac.uk](mailto:len@mcs.st-and.ac.uk)

Luís Matias  
Centro de Geofísica  
Universidade de Lisboa  
Lisbon, Portugal.

Award Number: N00014-11-1-0615  
Website: <http://www.creem.st-and.ac.uk/len/>

### **LONG-TERM GOALS**


Several of the current methods for density estimation of cetaceans using passive fixed acoustics rely on large, dense arrays of cabled hydrophones and/or auxiliary information from animal tagging projects conducted at the same time as the acoustic survey. Obtaining such data is costly, and may be impractical to the wider community interested in estimating cetacean density. Therefore, the goal of Cheap DECAF is to focus on the development of cetacean density estimation methods using sensors that are sparsely distributed and less expensive to deploy than the cabled military arrays focussed on to date.

### **OBJECTIVES**

Recordings of fin whales (*Balaenoptera physalus*) from a sparse array of Ocean Bottom Seismometers (OBS) will be the dataset used to develop and test a variety of density estimation methods. The OBS array was deployed for 1 year (2007-2008) off the south coast of Portugal, near the Straits of Gibraltar (Fig 1).

The specific objectives of the project are to:

1. demonstrate how cue-counting methods can be used efficiently to obtain estimates of density over long time periods and large spatial scales using directional sound sensors;
2. extend the methods to allow for uncertainty in the depth of vocalizing animals;
3. develop and apply methods based on tracking moving individual animals;
4. develop and apply methods based on measuring total sound energy in relevant frequency bands;
5. obtain baseline estimates of spatial density of fin whales in the study area.



**Fig. 1. Location of the array of 24 OBS sensors in the Atlantic off Portugal.**

## APPROACH

This project is in collaboration with Oregon State University (grant number: N00014-11-1-0606, PI: David Mellinger). The work is divided into 3 components, as follows:

Component 1: Fin whale vocalisations will be automatically detected and localised across the 1-year dataset, using existing methods. Established distance sampling methods using cue counting will be used to generate seasonal density estimates, and spatial patterns in density will be related to oceanographic features. Customised distance sampling software will be used (Thomas *et al.*, 2010). This component will also include the development of methods to account for the depth distribution of animals, which will involve a simulation exercise.

Component 2: This component will focus on estimating density where the unit of interest is the individual animal, rather than a cue, i.e., vocalisations. Methods to account for the movement of individual animals will be developed via a simulation study, building on work completed for a Master's thesis (DiTraglia, 2007).

Component 3: This component will develop a method that uses the total energy present in a species' frequency band as the statistic upon which a density estimate is made. The approach used will involve a Monte Carlo simulation and propagation modeling, to link density of animals to a given received energy level.

Components 1 and 2 is being led by the personnel involved with this project, and Component 3 is being led by Oregon State University. There is also a project management element, coordinating bi-monthly tele-conference progress meetings, and at least two face-to-face meetings, one in each project year.

## **WORK COMPLETED**

This project was due to start in April 2011, but this was delayed until September 2011 to allow the named post-doctoral research fellow, Danielle Harris, to complete her PhD.

## **RESULTS**

No results to date.

## **IMPACT/APPLICATIONS**

The main aim of Cheap DECAF is to make density estimation of cetaceans less costly and, therefore, more accessible to the wider scientific community. The methods developed here will be applicable to re-deployable arrays of both sea-bed mounted instruments (such as the OBS array) and surface buoys, so will hopefully increase our capability to monitor cetacean density in geographic areas of interest, including those where naval operations are conducted.

## **RELATED PROJECTS**

Cheap DECAF (Grant number: N00014-1-11-0606, PI: David Mellinger, Oregon State University)

DECAF: Density Estimation for Cetaceans from passive Acoustic Fixed sensors (ended February 2011)

## **REFERENCES**

DiTraglia, F.J. 2007. Models of Random Wildlife Movement with an Application to Distance Sampling. MMaths Statistics thesis, University of St Andrews.

Matias, L. 2009. Fin whale vocalizations recorded by the NEAREST OBS's: Preliminary evaluation. Unpublished.

Thomas, L., S.T. Buckland, E.A. Rexstad, J.L. Laake, S. Strindberg, S.L. Hedley, J.R.B. Bishop, T.A. Marques and K.P. Burnham. 2010. Distance software: design and analysis of distance sampling surveys for estimating population size. *J. Appl. Ecol.* 47:5-14.