

Characterization of Bed Morphodynamics Using Multibeam Echo Sounding (MBES) and Wavelet Transform (WT) Analysis

Marcelo H. Garcia
Ven Te Chow Hydrosystems Laboratory
Department of Civil and Environmental Engineering
University of Illinois at Urbana-Champaign
205 North Mathews Avenue
Urbana, Illinois 61801
Phone: (217) 244-4484, fax: (217) 333-0687, email: mhgarcia@illinois.edu

James L. Best
Department of Geology
University of Illinois at Urbana-Champaign
1301 W. Green St.
Urbana, Illinois 61801
Phone: (217) 244-1839, email: jimbest@illinois.edu

Award Number: N00014-11-1-0293
<http://www.vtchl.uiuc.edu/>

LONG-TERM GOALS

The overall objectives of this project are as follows:

Application of the Wavelet Transform (WT) technique in the spatial domain to characterize the spatial distribution of co-existing bedforms of multiple dimensions generated under different flow field scenarios (waves, unidirectional flows and combined flows).

Demonstrate the applicability of WT analysis for the description of bedforms frequently found in coastal and river environments. An improved description of bed features and their interactions will certainly contribute to a more complete understanding of theories describing bedform formation and evolution.

Improve overall understanding of small and large scale features, which is especially important because of their strong implications for the associated patterns of bed roughness characteristics, flow resistance and sediment transport.

Describe the hydrodynamic interaction of multiple sized bedforms and, conversely, the bed morphology given averaged hydrodynamic parameters; this approach could be used to develop and validate numerical approaches that will allow detailed examination of associated hydrodynamic and morphological conditions at inlets and river mouths.

OBJECTIVES

To extend the WT analysis technique developed by Cataño-Lopera *et al.* (2009) with laboratory observations, to the case of bed morphodynamics under field conditions at rivers and estuaries. For this purpose, the WT technique will be applied in the spatial domain rather than in the temporal domain, as conventionally used.

To apply the WT technique in the spatial domain to field measurements of bedforms in the Missouri River and the St. Clair River near Port Huron (USA-Canada). The bathymetric surveys employed a Multibeam Echo Sounding (MBES) system.

APPROACH

1- Starting from the 1D WT techniques by Cataño-Lopera *et al.* (2009), we developed a 2D whole-field technique based on the continuous wavelet transform method to analyze the bedform scales present in a particular bathymetry using the 2D Morlet wavelet function (Fig. 1).




Fig. 1. Morlet wavelet function for 1D space (left) and 2D space (right).

2- Calibration of the 2D WT tool using synthetic bathymetries generated numerically for simple bedform configurations.

3- Validation of the 2D WT tool using laboratory bathymetries from bedforms generated under standing waves.

4- Application of the 2D WT tool to bathymetric surveys from MBES swaths of two field sites: the Missouri River and the St. Clair River near Port Huron (USA-Canada).

The research team is composed by Prof. Marcelo H. Garcia as PI, Prof. James L. Best as co-PI, and Jose M. Mier as Ph.D. student. Development of the 2D WT tool was performed by Marcelo H. Garcia and Jose M. Mier. Data for validation was provided by Dr. Blake J. Landry from laboratory

experiments in the wave tank at the Ven Te Chow Hydrosystems Laboratory. The bathymetric surveys were performed by James L. Best who supplied the data for the field application of the technique.

WORK COMPLETED

1- The WT analysis tool for 2D bathymetry field has been developed. It is available as a MatLab® code that can be applied to an input bathymetry given as a ‘xyz’ file. It applies the 2D continuous wavelet transform technique using the 2D Morlet wavelet function for different scales and directions to the whole bathymetry field. It returns an output image of the original bathymetry with the different bedform scales present indicated by superimposed shadowed areas (Fig. 2). Additionally, the principal direction of bedform migration can be displayed by a superimposed arrow field.




Fig. 2. Results of the 2D WT tool applied to a unidirectional synthetic bathymetry field. Original bed elevation map (top) and identified bedform scales map (bottom).

2- The 2D WT has been calibrated using synthetic bathymetries both for scale (Fig. 2) and direction (Fig. 3) of the bedforms present.

3- The 2D WT tool has been validated using the bathymetric data sets of Landry (2011) which were generated under standing waves conditions. They include small scale bedforms (ripples) ranging from 5 cm to about 20 cm superimposed on top of large scale bedforms (bars) of about 4 m. The 2D WT tool has been able to identify the different bedform scales present and their location (Fig. 4).

4- The 2D WT tool has been applied to bathymetry data obtained from field surveys at the Missouri River and the St. Clair River near Port Huron (USA-Canada). Data was collected using a MBES system, which provided unrivalled resolution of the entire bottom morphology, with lateral coverage of each swath extending to nearly 5 times the flow depth. The x,y grid spacing of the datasets was 0.25 m and 0.5 m, respectively.




Fig. 3. Results of the 2D WT tool applied to a synthetic bathymetry field with different bedform scales and migration directions used during the calibration process. Original bed elevation map (left) and identified bedform scales and directions map (right).





Fig. 4. Results of the 2D WT tool applied to the laboratory bathymetry of Landry (2011) used for validation. Original bed elevation map (top) and identified bedform scales map (bottom).

From the analysis of the datasets, shadowed area maps were obtained showing the location, scale and direction of the dominant bedforms present. These results were overlaid on the original bathymetries to allow for easy identification of the features and their spatial distribution. In the Missouri River bathymetry (Fig. 5), two main scales were identified: the features on the bottom-right side of the image correspond to medium scale bedforms of 5 to 10 m in wavelength, while on the top-left of the image 20 m large scale bedforms were mapped by the 2D WT tool. Also, the arrows on the image help identify the direction of the bedforms, which most likely correspond to the direction of the dominant flow that contributed to generate each particular feature at that scale. Areas of superimposed bedforms were observed at other locations where multiple scales had been identified. In these cases, the

directions of the bedforms superimposing were rarely the same, suggesting the presence of different dominant flows for each scale.



Scales, dominant direction and location of bedforms based on 2D Wavelet Analysis




Fig. 5. Results of the 2D WT tool applied to the Missouri River bathymetry data. Original bed elevation map (top) and identified bedform scales map (bottom).

RESULTS

The 2D WT tool allows the analysis of bathymetric data sets to obtain dominant bedform scales, their location and migration direction for the whole field based on the continuous wavelet transform and the Morlet wavelet. This provides a significant improvement with respect the existing techniques which were only capable of 1D bedform scale analysis (Cataño-Lopera *et al.*, 2009).

The results are obtained as a shadowed area map showing the location, scale and direction of the dominant bedforms present superimposed on the original bathymetry. This allows scientists to easily identify the relevant bed features and their location, as well as their possible relations with the hydrodynamic conditions present.

Particularly interesting is the fact that the 2D WT tool is able to identify the existence of different bedform scales superimposed in the same region of the bathymetry field. The direction feature can be very useful in these cases to identify the dominant flows that led to the development of each one of the bedform scales.

IMPACT/APPLICATIONS

Wavelet analysis has been used in numerous studies in areas such as geophysics (Farge, 1992; Meyers et al., 1993; Weng and Lau, 1994; Liu, 1994; Gu and Philander, 1995; Massel, 2001), biology and medicine (Dettori and Semler, 2007), and acoustics (Lardies, 2007). However, this technique had not been extensively applied to the analysis of bedforms generated either under unidirectional currents, oscillatory flows or a combination of both, until now.

The interesting results obtained through this work under laboratory and field conditions suggest that the 2D WT tool has great potential for deployment in the field and could yield extremely useful information for naval operations.

RELATED PROJECTS

None

REFERENCES

- Best, J., Simmons, S., Parsons, D., Oberg, K., Czuba, J. and Malzone, C. (2010). "A new methodology for the quantitative visualization of coherent flow structures in alluvial channels using multiband echosounding (MBES)". *Geophys. Res. Lett.*, 37. LO6405, doi:10.1029/2009GL041852. [published, refereed].
- Cataño-Lopera, Y.A., Abad, J. D., and Garcia, M. H. (2009). "Characterization of bed form morphology generated under combined flows and currents using wavelet analysis". *Ocean Engineering*, vol. 36, issues 9-10, pp. 617-632.
- Cataño-Lopera, Y.A., and García, M.H. (2006a). "Geometry and Migration Characteristics of Bedforms under Waves and Currents: Part 1, Sandwave morphodynamics". *Coastal Engineering*, 53, 767-780. [published, refereed].

- Dettoni, L., and Semler, L. (2007). "A comparison of wavelet, ridgelet, and curvelet-based texture classification algorithms in computed tomography". *Computers in Biology and Medicine*, 37 (4), 486-498.
- Farge, M. (1992). "Wavelet transforms and their application to turbulence". *Annual Review of Fluid Mechanics*, 24, 395-457.
- Fedele, J.J. and M.H. Garcia (2005). "Roughness function for Alluvial Rivers with Dunes", in *Proceedings River, Coastal and Estuarine Morphodynamics*, RCEM, Parker & Garcia (Eds.), Urbana, Illinois. [published].
- Gu, D., and Philander, S.G.H. (1995). "Secular changes of annual and interannual variability in the Tropics during the past century". *J. of Climate*, 8, 864–876.
- Landry, B.J. (2011). "Sand bed morphodynamics under combined waves and vegetated conditions". Ph.D. Thesis, University of Illinois at Urbana-Champaign.
- Lardies, J. (2007). "Identification of a dynamical model for an acoustic enclosure using the wavelet Transform". *Applied Acoustics*, 68 (4), 473-490.
- Liu, P.C. (1994). "Wavelet spectrum analysis and ocean wind waves". *Wavelets in Geophysics*, E. Foufoula-Georgiou and P. Kumar, Eds., Academic Press, 151–166.
- Landry, B. J., Cataño-Lopera, Y. A., Hancock, M. J., Mei, C. C., and García, M. H. (2009). "Effect of Spatial Variation of a Wave Field on the Resulting Ripple Characteristics and Comparison to Present Ripple Predictors". *28th International Conference on Ocean, Offshore and Arctic Engineering*.
- Massel, S. R. (2001). "Wavelet analysis for processing of ocean surface wave records". *Ocean Engineering*, 28, 957–987.
- Meyers, S.D., Kelly, B.G., and O'Brien, J.J. (1993). "An introduction to wavelet analysis in oceanography and meteorology: With application to the dispersion of Yanai waves". *Mon. Wea. Rev.*, 121, 2858–2866.
- Parsons, D., Best, J.L., Hardy, R.J., Kostaschuk, R.A., Lane, S.N., and Orfeo, O. (2005). "The morphology and flow fields of three-dimensional dunes, Rio Paraná, Argentina: results from simultaneous multibeam echo sounding and acoustic Doppler current profiling". *Journal of Geophysical Research*, 110, F04S03, DOI:10.1029/2004JF000231.[published, refereed].
- Weng, H., Lau, K-M. (1994). "Wavelets, Period Doubling, and Time–Frequency Localization with Application to Organization of Convection over the Tropical Western Pacific". *J. Atmos. Sci.*, 51, 2523–2541.

PUBLICATIONS

- Czuba, Jonathan A.; Best, James L.; Oberg, Kevin A.; Parsons, Daniel R.; Jackson, P. Ryan; Garcia, Marcelo H.; Ashmore, P. (2011) "Bed morphology, flow structure, and sediment transport at the

outlet of Lake Huron and in the upper St. Clair River." *Journal of Great Lakes Research*, 37(3), 480-493. *Recognized with the IAGLR Chandler-Misener Award for most notable paper published in Journal of Great Lakes Research in 2011.*

García, M.H. (2008). Sediment transport and morphodynamics. Chapter 2 in ASCE Manual of Practice 110, *Sedimentation Engineering: Processes, Measurements, Modelling and Practice*. Edited by M. H. García, ASCE, Reston, Va.

Holmes, R.R. and Garcia, M.H. (2008). "Flow over Bedforms in a Large Sand-Bed River: A Field Investigation", *Journal of Hydraulic Research*, IAHR. Vol. 46, no. 3, pp. 322-333. [published, refereed].

Landry, B. J. and Garcia, M. H. (2007). "Bathymetric Evolution of a Sandy Bed under Transient Progressive Waves." *Proceedings of Coastal Sediments 2007*, 1: 2191-2198. [published, refereed]

Landry, B. J., Cataño-Lopera, Y. A., Hancock, M. J., Mei, C. C., and García, M. H. (2009). "Effect of Spatial Variation of a Wave Field on the Resulting Ripple Characteristics and Comparison to Present Ripple Predictors". *28th International Conference on Ocean, Offshore and Arctic Engineering*.

Mier, J. M., and Garcia, M.H. (2011). "Erosion of glacial till from the St. Clair River (Great Lakes basin)". *Journal of Great Lakes Research*, vol. 37, issue 3, pp. 399-410. [published, refereed].

Pedocchi, F., and García, M. H. (2009a). "Ripple Morphology under Oscillatory Flow. Part I: Prediction". *Journal of Geophysical Research*, vol. 114, C12014, 16 pp. [published, refereed].

Pedocchi, F., and García, M. H. (2009b). "Ripple Morphology under Oscillatory Flow. Part II: Experiments". *Journal of Geophysical Research*, vol. 114, C12015, 17 pp. [published, refereed].

Pedocchi, F., and García, M. H. (2011). "Acoustic measurement of suspended sediment concentration profiles in an oscillatory boundary layer". *Continental Shelf Research*, doi: 10.1016/j.csr.2011.05.013. [in press, refereed].

HONORS/AWARDS/PRIZES

Hunter Rouse Hydraulic Engineering Award, EWRI/ASCE, 2012.

Wesley Horner Award, best paper in *Journal of Environmental Engineering*, EWRI, ASCE, 2012.

International Association for Great Lakes Research Chandler-Misener Award for most notable paper published in *Journal of Great Lakes Research*, 2012.

National Award for Significant Contributions in Science and Technology, Panama Canal Authority, Government of Panama, April 2012.

Enrico Marchi Distinguished Lecture, Italian Association of Hydraulics, University of Florence, Italy, 2012.