

Origins of the Kuroshio and Mindanao Currents

Steven R. Jayne
Woods Hole Oceanographic Institution
MS 29, Clark 209A
Woods Hole, MA 02543-1541
phone: (508) 289-3520 fax: (508) 457-2163 email: sjayne@whoi.edu

Award Number: N00014-10-1-0268
<http://kirin.apl.washington.edu/okmc/>

LONG-TERM GOALS

The long-term goal of this work is to understand the ocean's circulation in the bifurcation region of the Kuroshio and Mindanao Currents. The westward flowing North Equatorial Current runs into the Philippine coast and bifurcates into the northward Kuroshio and the southward Mindanao Current. The partitioning of the flow into the Kuroshio and Mindanao Currents is an important observable. Quantifying these flows and understanding bifurcation dynamics are essential to improving predictions of regional circulation, and to characterizing property transports that ultimately affect Pacific climate.

OBJECTIVES

The objective of the OKMC program is to make observations of circulation in the area of the western Pacific Ocean where the North Equatorial Current runs into the western boundary near the Philippine Archipelago and bifurcates into the northward-flowing Kuroshio Current and the southward-flowing Mindanao Current. The observations made during OKMC include glider transects, surface drifters and profiling floats. It is anticipated that the mapped circulation will be used elucidate the ocean dynamics in the bifurcation region, and will help us understand what sets the partitioning of the current transports between the Kuroshio and Mindanao Currents. It will also provide improved estimates of the transport of the Mindanao Current, which has been poorly observed historically.

APPROACH

Our approach is to produce objective maps of the circulation utilizing the observations made by the surface drifters and profiling floats and combine them with the sea surface height (observed by satellite altimeters and referenced to the newest generation of geoids), and historical surface drifters and Argo float profiles and trajectories.

WORK COMPLETED

The objective mapping procedures have been tested and validated for the time-mean circulation. The mapping technique is different than traditional objective mapping techniques in that we impose physical constraints on the mapped circulation such as no flow through land in order to make it more consistent with the known physics.

RESULTS

Altimeter derived circulation is noisy, and unphysical near the coast. But estimates of the large-scale circulation from velocity (drifter) data alone are too weak due to noise in drifter data and the objective mapping technique. This can be overcome if we include both data types, sea surface height and surface drifter velocities to make a combined estimate. Additionally, the unrealistic flow through topography implied by the sea surface height can be overcome with a dynamic constraint that specifies no flow through topography. An objectively mapped surface circulation in the bifurcation region is shown in Figure 1.

We must also contend with the fact that the observations are not all simultaneous in time and space, and therefore need to be adjusted to the same time period. This can be accomplished by calculating the seasonal cycle and secular trend from the observations. For example, in Figure 2, we show the amplitudes of these terms. It is noteworthy that this region has the highest rate of sea level change anywhere in the world.




Figure 1: Objectively mapped time-averaged sea surface height (right) and surface geostrophic current (left). The mapped fields are estimated by objectively mapping the sea surface height, surface drifters (corrected for Ekman velocities) and the constraint of no flow through topography.




Figure 2: Annual amplitude of the annual cycle (right) and secular trend (left) in sea surface height. Note this region has the largest secular trend of increasing sea surface in the world.

IMPACT/APPLICATIONS

It is anticipated that the mapped circulation will be used elucidate the ocean dynamics in the bifurcation region, and will help us understand what sets the partitioning of the current transports between the Kuroshio and Mindanao Currents. It will also provide improved estimates of the transport of the Mindanao Current which has been poorly observed historically.

RELATED PROJECTS

Similar maps of the sea surface height and velocity field are being developed for the global ocean for the Argo program (<http://www.argo.net>) and the Ocean Surface Topography Science team (<https://sealevel.jpl.nasa.gov/science/ostscienceteam/>).